

ZU-TH 12/05
hep-ph/0507061
July 2005

Two-Loop Quark and Gluon Form Factors in Dimensional Regularisation

T. Gehrmann, T. Huber and D. Maître

*Institut für Theoretische Physik, Universität Zürich, Winterthurerstrasse 190,
CH-8057 Zürich, Switzerland*

Abstract

We compute the two-loop corrections to the massless quark form factor $\gamma^* \rightarrow q\bar{q}$ and gluon form factor $H \rightarrow gg$ to all orders in the dimensional regularisation parameter $\epsilon = (4 - d)/2$. The two-loop contributions to the form factors are reduced to linear combinations of master integrals, which are computed in a closed form, expressed as Γ -functions and generalised hypergeometric functions of unit argument. Using the newly developed **HypExp**-package, these can be expanded to any desired order, yielding Laurent expansions in ϵ . We provide expansions of the form factors to order ϵ^2 , as required for ultraviolet renormalisation and infrared factorisation of the three-loop form factors.

1 Introduction

The infrared pole structure of renormalised multi-loop amplitudes in dimensional regularisation with $d = 4 - 2\epsilon$ space-time dimensions can be predicted from an infrared factorisation formula, which was first conjectured in [1], where it was formulated up to two loops. A proof of the formula, together with an explicit formulation up to three loops was derived later in [2]. The simplest multi-loop amplitudes where the infrared factorisation formula can be applied are three-point functions, involving two partons coupled to an external current: the quark form factor $\gamma^* \rightarrow q\bar{q}$ and the gluon form factor $H \rightarrow gg$. The QCD corrections to these form factors can in particular be used to fix a priori unknown constants in the infrared factorisation formula, thus enabling an unambiguous prediction for multi-loop amplitudes involving more than two external partons.

In the infrared factorisation formula for a given form factor (or more generally for a given multi-leg amplitude) at a certain number of loops, infrared singularity operators act on the form factor evaluated with a lower number of loops. The infrared singularity operators contain explicit infrared poles $1/\epsilon^2$ and $1/\epsilon$. They do therefore project subleading terms in ϵ from the lower order form factors.

At present, two-loop corrections to the massless quark [3] and gluon [4] form factors are known to order ϵ^0 . Two-loop corrections to this order were also obtained for massive quarks [5]. The infrared structure of the massless form factors and infrared cancellations with real radiation contributions are described in detail in [6]. Very recently, results to order ϵ^2 were obtained for the quark form factor [7].

The calculation of these corrections proceeds through a reduction [8–11] of all two-loop Feynman integrals appearing in the form factors to a small set of master integrals. The reduction is exact in ϵ , such that the evaluation of the form factors is limited only by the order to which the master integrals can be computed. The massless two-loop form factors contain three two-loop master integrals, which can be computed either using various analytical methods [12] or numerically order-by-order in their Laurent expansion using the sector decomposition algorithm [13]. Up to now, exact expressions were known only for two of these master integrals, while the third (the so-called two-loop crossed triangle graph) was known only as a Laurent expansion up to finite terms [14].

In this letter, we derive an exact expression for the two-loop crossed triangle graph in terms of generalised hypergeometric functions of unit argument in Section 2. Using the **HypExp**-package [15] for the Laurent expansion of generalised hypergeometric functions, this can be expanded to any desired order in ϵ . Together with the exact expressions for the one- and two-loop quark and gluon form factors in Section 3, this allows the expansion of these form factors to higher orders in ϵ . For illustration, we list the one-loop form factors to order ϵ^4 and the two-loop form factors to order ϵ^2 in Section 4; these orders appear for example in the infrared factorisation of the corresponding three-loop form factors. Finally, Section 5 contains conclusions and an outlook.

2 Two-loop master integrals

The virtual two-loop vertex master integrals were first derived to order ϵ^0 in [14] in the context of the calculation of the two-loop quark form factor [3]. All but the crossed triangle graph A_6 can be expressed in terms of Γ -functions to all orders in ϵ .

Factoring out a common

$$S_\Gamma = \left(\frac{(4\pi)^\epsilon}{16\pi^2 \Gamma(1-\epsilon)} \right), \quad (1)$$

and introducing $q^2 = (p_1 + p_2)^2$, they read

$$\begin{aligned} A_{2,\text{LO}} &= \text{Diagram: a circle with an incoming line from the left labeled } p_{12} \text{ and an outgoing line to the right.} \\ &= \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^2 (k - p_1 - p_2)^2} \\ &= S_\Gamma (-q^2)^{-\epsilon} \frac{\Gamma(1+\epsilon) \Gamma^3(1-\epsilon)}{\Gamma(2-2\epsilon)} \frac{i}{\epsilon}, \end{aligned} \quad (2)$$

$$\begin{aligned} A_3 &= \text{Diagram: a circle with an incoming line from the left labeled } p_{12} \text{ and an outgoing line to the right labeled } p_1. \\ &= \int \frac{d^d k}{(2\pi)^d} \int \frac{d^d l}{(2\pi)^d} \frac{1}{k^2 l^2 (k - l - p_1 - p_2)^2} \\ &= S_\Gamma^2 (-q^2)^{1-2\epsilon} \frac{\Gamma(1+2\epsilon) \Gamma^5(1-\epsilon)}{\Gamma(3-3\epsilon)} \frac{-1}{2(1-2\epsilon)\epsilon}, \end{aligned} \quad (3)$$

$$\begin{aligned} A_4 &= \text{Diagram: a circle with an incoming line from the left labeled } p_{12} \text{ and two outgoing lines to the right labeled } p_1 \text{ and } p_2. \\ &= \int \frac{d^d k}{(2\pi)^d} \int \frac{d^d l}{(2\pi)^d} \frac{1}{k^2 l^2 (k - p_1 - p_2)^2 (k - l - p_1)^2} \\ &= S_\Gamma^2 (-q^2)^{-2\epsilon} \frac{\Gamma(1-2\epsilon) \Gamma(1+\epsilon) \Gamma^4(1-\epsilon) \Gamma(1+2\epsilon)}{\Gamma(2-3\epsilon)} \frac{-1}{2(1-2\epsilon)\epsilon^2}. \end{aligned} \quad (4)$$

No exact expression for A_6 was known up to now. Following the steps outlined in [14], we obtain

$$\begin{aligned} A_6 &= \text{Diagram: a crossed triangle graph with an incoming line from the left labeled } p_{12} \text{ and two outgoing lines to the right labeled } p_1 \text{ and } p_2. \\ &= \int \frac{d^d k}{(2\pi)^d} \int \frac{d^d l}{(2\pi)^d} \frac{1}{k^2 l^2 (k - p_1 - p_2)^2 (k - l)^2 (k - l - p_2)^2 (l - p_1)^2} \\ &= S_\Gamma^2 (-q^2)^{-2-2\epsilon} \left[- \frac{\Gamma^3(1-\epsilon) \Gamma(1+\epsilon) \Gamma^4(1-2\epsilon) \Gamma^3(1+2\epsilon)}{\epsilon^4 \Gamma^2(1-4\epsilon) \Gamma(1+4\epsilon)} \right. \\ &\quad \left. + \frac{\Gamma^4(1-\epsilon) \Gamma(1+\epsilon) \Gamma(1-2\epsilon) \Gamma(1+2\epsilon)}{2\epsilon^4 \Gamma(1-3\epsilon)} {}_3F_2(1, -4\epsilon, -2\epsilon; 1-3\epsilon, 1-2\epsilon; 1) \right] \end{aligned}$$

$$\begin{aligned} & - \frac{4\Gamma^4(1-\epsilon)\Gamma(1-2\epsilon)\Gamma(1+2\epsilon)}{\epsilon^2(1+\epsilon)(1+2\epsilon)\Gamma(1-4\epsilon)} {}_3F_2(1, 1, 1+2\epsilon; 2+\epsilon, 2+2\epsilon; 1) \\ & - \frac{\Gamma^5(1-\epsilon)\Gamma(1+2\epsilon)}{2\epsilon^4\Gamma(1-3\epsilon)} {}_4F_3(1, 1-\epsilon, -4\epsilon, -2\epsilon; 1-3\epsilon, 1-2\epsilon, 1-2\epsilon; 1) \Big] . \end{aligned} \quad (5)$$

While $A_{2,LO}$, A_3 and A_4 can be expanded using any standard computer algebra programme, the expansion of A_6 requires the expansion of generalised hypergeometric functions in their parameters. For this purpose, a dedicated package, **HypExp** [15], was developed recently. Using this, we obtain the eighth-order expansion:

$$\begin{aligned} A_6 = & S_F^2 (-q^2)^{-2-2\epsilon} \left[-\frac{1}{\epsilon^4} + \frac{5\pi^2}{6\epsilon^2} + \frac{27}{\epsilon}\zeta_3 + \frac{23\pi^4}{36} + (117\zeta_5 - 8\pi^2\zeta_3)\epsilon \right. \\ & + \left(\frac{19\pi^6}{315} - 267\zeta_3^2 \right) \epsilon^2 + \left(-\frac{109\pi^4}{10}\zeta_3 - 40\pi^2\zeta_5 - 6\zeta_7 \right) \epsilon^3 \\ & \left. + \left(44\pi^2\zeta_3^2 - \frac{1073\pi^8}{3024} - 2466\zeta_3\zeta_5 + 264\zeta_{5,3} \right) \epsilon^4 + \mathcal{O}(\epsilon^5) \right] , \end{aligned} \quad (6)$$

where we encountered a multiple zeta value in the last term.

3 Quark and gluon form factors at two loops

The tree-level quark and gluon form factors are obtained by normalising the corresponding tree-level vertex functions to unity:

$$F_q^{(0l)} = 1, \quad F_g^{(0l)} = 1. \quad (7)$$

The unrenormalised one-loop and two-loop form factors are calculated from the relevant Feynman diagrams. Using integration-by-parts [8] and Lorentz invariance [10] identities (which can be solved symbolically for massless two-loop vertex integrals, see the appendix of [16]), these can be reduced [9–11] to the master integrals listed in Section 2.

The unrenormalised one-loop quark and gluon form factors read:

$$F_q^{(1l,B)} = -ig^2 \frac{N^2 - 1}{N} \frac{d^2 - 7d + 16}{2(d-4)} A_{2,LO}, \quad (8)$$

$$F_g^{(1l,B)} = ig^2 N \frac{d^3 - 16d^2 + 68d - 88}{(d-4)(d-2)} A_{2,LO}, \quad (9)$$

where $N = 3$ is the number of colours and g is the bare QCD coupling parameter.

The unrenormalised two-loop quark and gluon form factors for N_F massless quark flavours are:

$$\begin{aligned} F_q^{(2l,B)} = & g^4 \frac{N^2 - 1}{N} \left\{ -\frac{N^2 - 1}{N} \frac{(d^2 - 7d + 16)^2}{4(d-4)^2} A_{2,LO}^2 \right. \\ & \left. + N \frac{(d^5 - 18d^4 + 138d^3 - 552d^2 + 1144d - 980)(3d-8)}{2(d-3)(d-4)^3} \frac{A_3}{q^2} \right\} \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{N} \frac{(9d^6 - 358d^5 + 4309d^4 - 24466d^3 + 72896d^2 - 110064d + 66080)(3d-8)}{16(d-3)(d-4)^3(2d-7)} \frac{A_3}{q^2} \\
& + N \frac{3d^6 - 82d^5 + 819d^4 - 4030d^3 + 10344d^2 - 12824d + 5632}{4(d-1)(d-4)^2(3d-8)} A_4 \\
& - \frac{1}{N} \frac{(21d^6 - 789d^5 + 9422d^4 - 53864d^3 + 163200d^2 - 253472d + 159232)}{16(3d-8)(2d-7)(d-4)^2} A_4 \\
& + N_F \frac{(3d^3 - 31d^2 + 110d - 128)(d-2)}{2(d-1)(d-4)(3d-8)} A_4 \\
& - \frac{1}{N} \frac{d^3 - 20d^2 + 104d - 176}{32(2d-7)} (q^2)^2 A_6 \Big\} , \tag{10} \\
F_g^{(2l,B)} = g^4 \Big\{ & - N^2 \frac{(d^3 - 16d^2 + 68d - 88)^2}{(d-4)^2(d-2)^2} A_{2,\text{LO}}^2 \\
& - N^2 \frac{1}{2(d-1)(d-2)^2(d-3)(d-4)^3(2d-5)(2d-7)} \Big(192d^{10} - 6947d^9 \\
& + 105470d^8 - 907248d^7 + 4958664d^6 - 18113645d^5 + 44930982d^4 \\
& - 74791460d^3 + 79854504d^2 - 49204128d + 13194496 \Big) \frac{A_3}{q^2} \\
& - N N_F \frac{2d^6 - 45d^5 + 377d^4 - 1610d^3 + 3868d^2 - 5136d + 3008}{(d-1)(d-2)(d-3)(d-4)^2} \frac{A_3}{q^2} \\
& + \frac{N_F}{N} \frac{1}{4(d-2)(d-3)(d-4)^2(2d-5)(2d-7)} \Big(70d^7 - 1663d^6 + 16290d^5 \\
& - 86031d^4 + 266004d^3 - 483356d^2 + 479360d - 200704 \Big) \frac{A_3}{q^2} \\
& - N^2 \frac{1}{2(d-1)(d-2)^2(d-4)^2(2d-5)(2d-7)} \Big(108d^8 - 2661d^7 + 28822d^6 \\
& - 177546d^5 + 674735d^4 - 1607602d^3 + 2325996d^2 - 1848920d + 607968 \Big) A_4 \\
& + N N_F \frac{2d^4 - 28d^3 + 130d^2 - 228d + 104}{(d-1)(d-2)(d-4)} A_4 \\
& - \frac{N_F}{N} \frac{(46d^4 - 545d^3 + 2395d^2 - 4606d + 3248)(d-6)}{4(d-2)(d-4)(2d-5)(2d-7)} A_4 \\
& - N^2 \frac{3(3d-8)(d-3)}{4(2d-5)(2d-7)} (q^2)^2 A_6 \\
& - \frac{N_F}{N} \frac{(d-4)(2d^3 - 25d^2 + 94d - 112)}{8(d-2)(2d-5)(2d-7)} (q^2)^2 A_6 \Big\} . \tag{11}
\end{aligned}$$

The renormalised form factors are obtained by introducing the renormalised QCD coupling constant and the renormalised effective coupling of H to the gluon field strength [4], and subsequent expansion in powers of the renormalised coupling.

4 Expansion of two-loop form factors

The renormalised form factors are expanded in the renormalised coupling constant. In the $\overline{\text{MS}}$ scheme, the bare coupling $\alpha_0 = g^2/(4\pi)$ is related to the renormalised coupling $\alpha_s \equiv \alpha_s(\mu^2)$, evaluated at the renormalisation scale μ^2 by

$$\alpha_0 \mu_0^{2\epsilon} S_\epsilon = \alpha_s \mu^{2\epsilon} \left[1 - \frac{11N - 2N_F}{6\epsilon} \left(\frac{\alpha_s}{2\pi} \right) + \mathcal{O}(\alpha_s^2) \right], \quad (12)$$

where

$$S_\epsilon = (4\pi)^\epsilon e^{-\epsilon\gamma} \quad \text{with the Euler constant } \gamma = 0.5772\dots$$

and μ_0^2 is the mass parameter introduced in dimensional regularisation to maintain a dimensionless coupling in the bare QCD Lagrangian density. For simplicity, we set $\mu^2 = q^2$. If the squared momentum transfer q^2 is space-like ($q^2 < 0$), the form factors are real, while they acquire imaginary parts for time-like q^2 . These imaginary parts (and corresponding real parts) arise from the ϵ -expansion of

$$\Delta(q^2) = (-\text{sgn}(q^2) - i0)^{-\epsilon}. \quad (13)$$

The renormalised form factors can then be written as

$$F_{q,g}(q^2) = 1 + \left(\frac{\alpha_s}{2\pi} \Delta(q^2) \right) F_{q,g}^{(1)} + \left(\frac{\alpha_s}{2\pi} \Delta(q^2) \right)^2 F_{q,g}^{(2)} + \mathcal{O}(\alpha_s^3). \quad (14)$$

Expanding the first and second order coefficients of the form factors to ϵ^4 and ϵ^2 respectively, we obtain:

$$\begin{aligned} F_q^{(1)} &= \left(N - \frac{1}{N} \right) \left[-\frac{1}{2\epsilon^2} - \frac{3}{4\epsilon} - 2 + \frac{\pi^2}{24} + \left(-4 + \frac{\pi^2}{16} + \frac{7}{6}\zeta_3 \right) \epsilon \right. \\ &\quad + \left(-8 + \frac{\pi^2}{6} + \frac{7}{4}\zeta_3 - \frac{47\pi^4}{2880} \right) \epsilon^2 \\ &\quad + \left(-16 - \frac{7\pi^2}{72}\zeta_3 + \frac{14}{3}\zeta_3 + \frac{31}{10}\zeta_5 + \frac{\pi^2}{3} + \frac{47\pi^4}{1920} \right) \epsilon^3 \\ &\quad \left. + \left(-32 - \frac{7\pi^2}{48}\zeta_3 + \frac{28}{3}\zeta_3 - \frac{49}{36}\zeta_3^2 + \frac{93}{20}\zeta_5 + \frac{2\pi^2}{3} + \frac{47\pi^4}{720} + \frac{949\pi^6}{241920} \right) \epsilon^4 \right] + \mathcal{O}(\epsilon^5), \quad (15) \end{aligned}$$

$$\begin{aligned} F_g^{(1)} &= N \left[-\frac{1}{\epsilon^2} - \frac{11}{6\epsilon} + \frac{\pi^2}{12} + \left(-1 + \frac{7}{3}\zeta_3 \right) \epsilon + \left(-3 + \frac{47\pi^4}{1440} \right) \epsilon^2 \right. \\ &\quad + \left(-7 - \frac{7\pi^2}{36}\zeta_3 + \frac{31}{5}\zeta_5 + \frac{\pi^2}{12} \right) \epsilon^3 \\ &\quad \left. + \left(-15 + \frac{7}{3}\zeta_3 - \frac{49}{18}\zeta_3^2 + \frac{\pi^2}{4} + \frac{949\pi^6}{120960} \right) \epsilon^4 \right] + \frac{N_F}{3\epsilon} + \mathcal{O}(\epsilon^5), \quad (16) \end{aligned}$$

$$F_q^{(2)} = \left(N - \frac{1}{N} \right) \left\{ N \left[\frac{1}{8\epsilon^4} + \frac{17}{16\epsilon^3} + \frac{433}{288\epsilon^2} \right. \right.$$

$$\begin{aligned}
& + \frac{1}{\epsilon} \left(\frac{4045}{1728} - \frac{11\pi^2}{96} + \frac{7}{24}\zeta_3 \right) + \left(-\frac{9083}{10368} - \frac{521\pi^2}{1728} + \frac{13}{18}\zeta_3 + \frac{23\pi^4}{2880} \right) \\
& + \left(-\frac{1244339}{62208} - \frac{11\pi^2}{48}\zeta_3 + \frac{4235}{432}\zeta_3 + \frac{163}{40}\zeta_5 - \frac{10427\pi^2}{10368} + \frac{29\pi^4}{1440} \right) \epsilon \\
& + \left(-\frac{36528395}{373248} - \frac{77\pi^2}{432}\zeta_3 + \frac{109019}{2592}\zeta_3 - \frac{403}{72}\zeta_3^2 + \frac{529}{30}\zeta_5 \right. \\
& \quad \left. - \frac{181451\pi^2}{62208} + \frac{8759\pi^4}{51840} + \frac{47\pi^6}{7560} \right) \epsilon^2 \Big] \\
& + \frac{1}{N} \left[-\frac{1}{8\epsilon^4} - \frac{3}{8\epsilon^3} + \frac{1}{\epsilon^2} \left(-\frac{41}{32} + \frac{\pi^2}{48} \right) \right. \\
& + \frac{1}{\epsilon} \left(-\frac{221}{64} + \frac{4}{3}\zeta_3 \right) + \left(-\frac{1151}{128} - \frac{17\pi^2}{192} + \frac{29}{8}\zeta_3 + \frac{13\pi^4}{576} \right) \\
& + \left(-\frac{5741}{256} - \frac{7\pi^2}{18}\zeta_3 + \frac{839}{48}\zeta_3 + \frac{23}{10}\zeta_5 - \frac{71\pi^2}{128} + \frac{19\pi^4}{320} \right) \epsilon \\
& + \left(-\frac{27911}{512} - \frac{9\pi^2}{16}\zeta_3 + \frac{6989}{96}\zeta_3 - \frac{163}{9}\zeta_3^2 + \frac{231}{40}\zeta_5 - \frac{613\pi^2}{256} + \frac{3401\pi^4}{11520} - \frac{223\pi^6}{17280} \right) \epsilon^2 \Big] \\
& + N_F \left[-\frac{1}{8\epsilon^3} - \frac{1}{18\epsilon^2} + \frac{1}{\epsilon} \left(\frac{65}{432} + \frac{\pi^2}{48} \right) + \left(\frac{4085}{2592} + \frac{23\pi^2}{432} + \frac{1}{36}\zeta_3 \right) \right. \\
& + \left(\frac{108653}{15552} - \frac{119}{108}\zeta_3 + \frac{497\pi^2}{2592} + \frac{\pi^4}{1440} \right) \epsilon \\
& + \left. \left(\frac{2379989}{93312} - \frac{5\pi^2}{54}\zeta_3 - \frac{3581}{648}\zeta_3 - \frac{59}{60}\zeta_5 + \frac{9269\pi^2}{15552} - \frac{145\pi^4}{10368} \right) \epsilon^2 \right] \Big\} + \mathcal{O}(\epsilon^3), \quad (17)
\end{aligned}$$

$$\begin{aligned}
F_g^{(2)} &= N^2 \left[\frac{1}{2\epsilon^4} + \frac{77}{24\epsilon^3} + \frac{1}{\epsilon^2} \left(\frac{175}{72} - \frac{\pi^2}{24} \right) \right. \\
& + \frac{1}{\epsilon} \left(-\frac{119}{54} - \frac{25}{12}\zeta_3 - \frac{11\pi^2}{144} \right) + \left(\frac{8237}{648} - \frac{33}{4}\zeta_3 + \frac{67\pi^2}{144} - \frac{7\pi^4}{240} \right) \\
& + \left(\frac{200969}{3888} + \frac{23\pi^2}{72}\zeta_3 - \frac{1139}{108}\zeta_3 + \frac{71}{20}\zeta_5 + \frac{53\pi^2}{108} - \frac{1111\pi^4}{8640} \right) \epsilon \\
& + \left. \left(\frac{4082945}{23328} - \frac{11\pi^2}{216}\zeta_3 - \frac{13109}{162}\zeta_3 + \frac{901}{36}\zeta_3^2 - \frac{341}{20}\zeta_5 + \frac{85\pi^2}{1296} - \frac{1943\pi^4}{8640} + \frac{257\pi^6}{6720} \right) \epsilon^2 \right] \\
& + N N_F \left[-\frac{7}{12\epsilon^3} - \frac{13}{12\epsilon^2} + \frac{1}{\epsilon} \left(\frac{155}{216} + \frac{\pi^2}{72} \right) \right. \\
& + \left. \left(-\frac{5905}{1296} + \frac{1}{2}\zeta_3 - \frac{5\pi^2}{72} \right) + \left(-\frac{162805}{7776} - \frac{95}{54}\zeta_3 - \frac{11\pi^2}{432} + \frac{7\pi^4}{1440} \right) \epsilon \right]
\end{aligned}$$

$$\begin{aligned}
& + \left(-\frac{3663205}{46656} + \frac{31\pi^2}{108}\zeta_3 + \frac{274}{81}\zeta_3 - \frac{9}{10}\zeta_5 + \frac{883\pi^2}{2592} - \frac{73\pi^4}{2592} \right) \epsilon^2 \Big] \\
& + \frac{N_F}{N} \left[-\frac{1}{8\epsilon} + \left(\frac{67}{48} - \zeta_3 \right) + \left(\frac{2027}{288} - \frac{23}{6}\zeta_3 - \frac{7\pi^2}{144} - \frac{\pi^4}{54} \right) \epsilon \right. \\
& \left. + \left(\frac{47491}{1728} + \frac{5\pi^2}{18}\zeta_3 - \frac{281}{18}\zeta_3 - 4\zeta_5 - \frac{209\pi^2}{864} - \frac{23\pi^4}{324} \right) \epsilon^2 \right] + N_F^2 \frac{1}{9\epsilon^2} + \mathcal{O}(\epsilon^3) . \quad (18)
\end{aligned}$$

5 Conclusions and outlook

In this letter, we computed the two-loop quark and gluon form factors to all orders in the dimensional regularisation parameter ϵ . The principal ingredient to this calculation is the two-loop crossed triangle graph A_6 , for which we computed an exact expression in terms of generalised hypergeometric functions of unit argument, which can be expanded to any desired order in ϵ using the **HypExp**-package.

A potential application of the form factors derived here is the extraction of the complete set of infrared pole terms of the genuine three-loop quark form factor from the recently derived three-loop splitting and coefficient functions in deep inelastic scattering [17]. In turn, these allow to fix the yet unknown hard radiation constants in the infrared factorisation formula at three loops. Parts of these constants were derived previously from $\mathcal{N} = 4$ supersymmetry relations [18, 19].

The two-loop vertex master integrals feature as subtopologies in the reduction of the three-loop form factor contributions, appearing if one of the three loops is disconnected from the others by pinching the connecting propagators. In this case, their terms to ϵ^2 are required.

The calculation presented here illustrates the applicability of the **HypExp**-package in the calculation of multi-loop corrections in quantum field theory. Functions similar to those which were expanded here appear also in multi-particle phase space integrals in massless [20] and massive decay processes [21]. Since all these integrals correspond to particular cuts of multi-loop two-point functions, one might expect that three-loop and four-loop two-point functions could also be expanded using **HypExp** to high orders in ϵ , as required for multi-loop calculations of fully inclusive observables [22].

Note added: While finalising this letter, an independent paper addressing very similar issues appeared. In hep-ph/0507039 [7], Moch, Vermaseren and Vogt compute the two-loop quark form factor to order ϵ^2 and apply it in the extraction of the pole parts of the three-loop quark form factor from deep inelastic coefficient functions. In this paper, the hard radiation constants for infrared factorisation at three-loops and related resummation coefficients are extracted for processes involving quarks only. Expanding our unrenormalised quark form factor (10) to order ϵ^2 , we confirm the result (B.1) of [7].

Acknowledgements

We wish to thank Gudrun Heinrich for independent numerical checks of the **HypExp**-Laurent expansions using the sector decomposition method described in [13].

This research was supported by the Swiss National Science Foundation (SNF) under contract 200021-101874.

References

- [1] S. Catani, Phys. Lett. B **427** (1998) 161 [hep-ph/9802439].
- [2] G. Sterman and M.E. Tejeda-Yeomans, Phys. Lett. B **552** (2003) 48 [hep-ph/0210130].
- [3] G. Kramer and B. Lampe, Z. Phys. C **34** (1987) 497; **42** (1989) 504(E);
T. Matsuura and W.L. van Neerven, Z. Phys. C **38** (1988) 623;
T. Matsuura, S.C. van der Maarck and W.L. van Neerven, Nucl. Phys. B **319** (1989) 570.
- [4] R.V. Harlander, Phys. Lett. B **492** (2000) 74 [hep-ph/0007289].
- [5] W. Bernreuther *et al.*, Nucl. Phys. B **706** (2005) 245 [hep-ph/0406046]; **712** (2005) 229 [hep-ph/0412259]; hep-ph/0504190.
- [6] A. Gehrmann-De Ridder, T. Gehrmann and E.W.N. Glover, Nucl. Phys. B **691** (2004) 195 [hep-ph/0403057]; Phys. Lett. B **612** (2005) 36 [hep-ph/0501291]; **612** (2005) 49 [hep-ph/0502110]; hep-ph/0505111.
- [7] S. Moch, J.A.M. Vermaseren and A. Vogt, hep-ph/0507039.
- [8] F.V. Tkachov, Phys. Lett. **100B** (1981) 65;
K.G. Chetyrkin and F.V. Tkachov, Nucl. Phys. **B192** (1981) 159.
- [9] S. Laporta, Int. J. Mod. Phys. A **15** (2000) 5087 [hep-ph/0102033].
- [10] T. Gehrmann and E. Remiddi, Nucl. Phys. B **580** (2000) 485 [hep-ph/9912329].
- [11] C. Anastasiou and A. Lazopoulos, JHEP **0407** (2004) 046 [hep-ph/0404258].
- [12] V.A. Smirnov, *Evaluating Feynman Integrals*, Springer Tracts of Modern Physics (Heidelberg, 2004).
- [13] T. Binoth and G. Heinrich, Nucl. Phys. B **585** (2000) 741 [hep-ph/0004013]; **680** (2004) 375 [hep-ph/0305234].
- [14] R.J. Gonsalves, Phys. Rev. **D28** (1983) 1542;
W.L. van Neerven, Nucl. Phys. B **268** (1986) 453;
G. Kramer and B. Lampe, J. Math. Phys. **28** (1987) 945.
- [15] T. Huber and D. Maître, <http://www-theorie.physik.unizh.ch/~maitreda/HypExp>, hep-ph/0507094.
- [16] C. Anastasiou, T. Gehrmann, C. Oleari, E. Remiddi and J.B. Tausk, Nucl. Phys. B **580** (2000) 577 [hep-ph/0003261].

- [17] S. Moch, J.A.M. Vermaseren and A. Vogt, Nucl. Phys. B **688** (2004) 101 [hep-ph/0403192]; **691** (2004) 129 [hep-ph/0404111]; hep-ph/0504242; hep-ph/0506288.
- [18] A.V. Kotikov, L.N. Lipatov, A.I. Onishchenko and V.N. Velizhanin, Phys. Lett. B **595** (2004) 521 [hep-th/0404092].
- [19] Z. Bern, L.J. Dixon and V.A. Smirnov, hep-th/0505205.
- [20] A. Gehrmann-De Ridder, T. Gehrmann and G. Heinrich, Nucl. Phys. B **682** (2004) 265 [hep-ph/0311276].
- [21] T. Huber, E. Lunghi, M. Misiak and D. Wyler, in preparation.
- [22] S. Bekavac, hep-ph/0505174.